

At this value of  $x$ ,

$$y = 9 - \frac{9}{2} = \frac{9}{2} \quad \text{and} \quad f(x, y) = f\left(\frac{9}{2}, \frac{9}{2}\right) = -\frac{41}{2}.$$

**Summary.** We list all the candidates: 4, 2,  $-61$ , 3,  $-(41/2)$ . The maximum is 4, which  $f$  assumes at (1, 1). The minimum is  $-61$ , which  $f$  assumes at (0, 9) and (9, 0).  $\square$

## Conclusion

Despite the power of Theorem 7, we urge you to remember its limitations. It does not apply to boundary points of a function's domain, where it is possible for a function to have extreme values along with nonzero derivatives. And it does not apply to points where either  $f_x$  or  $f_y$  fails to exist.

### Summary of Max-Min Tests

The extreme values of  $f(x, y)$  can occur only at

- (i) **boundary points** of the domain of  $f$ ,
- (ii) **critical points** (interior points where  $f_x = f_y = 0$  or points where  $f_x$  or  $f_y$  fail to exist).

If the first and second order partial derivatives of  $f$  are continuous throughout a disk centered at a point  $(a, b)$ , and  $f_x(a, b) = f_y(a, b) = 0$ , you may be able to classify  $f(a, b)$  with the **second derivative test**:

- (i)  $f_{xx} < 0$  and  $f_{xx}f_{yy} - f_{xy}^2 > 0$  at  $(a, b) \Rightarrow$  **local maximum**,
- (ii)  $f_{xx} > 0$  and  $f_{xx}f_{yy} - f_{xy}^2 > 0$  at  $(a, b) \Rightarrow$  **local minimum**,
- (iii)  $f_{xx}f_{yy} - f_{xy}^2 < 0$  at  $(a, b) \Rightarrow$  **saddle point**,
- (iv)  $f_{xx}f_{yy} - f_{xy}^2 = 0$  at  $(a, b) \Rightarrow$  **test is inconclusive**.

## Exercises 12.8

### Finding Local Extrema

Find all the local maxima, local minima, and saddle points of the functions in Exercises 1–30.

1.  $f(x, y) = x^2 + xy + y^2 + 3x - 3y + 4$
2.  $f(x, y) = x^2 + 3xy + 3y^2 - 6x + 3y - 6$
3.  $f(x, y) = 2xy - 5x^2 - 2y^2 + 4x + 4y - 4$
4.  $f(x, y) = 2xy - 5x^2 - 2y^2 + 4x - 4$
5.  $f(x, y) = x^2 + xy + 3x + 2y + 5$
6.  $f(x, y) = y^2 + xy - 2x - 2y + 2$
7.  $f(x, y) = 5xy - 7x^2 + 3x - 6y + 2$
8.  $f(x, y) = 2xy - x^2 - 2y^2 + 3x + 4$
9.  $f(x, y) = x^2 - 4xy + y^2 + 6y + 2$


10.  $f(x, y) = 3x^2 + 6xy + 7y^2 - 2x + 4y$
11.  $f(x, y) = 2x^2 + 3xy + 4y^2 - 5x + 2y$
12.  $f(x, y) = 4x^2 - 6xy + 5y^2 - 20x + 26y$
13.  $f(x, y) = x^2 - y^2 - 2x + 4y + 6$
14.  $f(x, y) = x^2 - 2xy + 2y^2 - 2x + 2y + 1$
15.  $f(x, y) = x^2 + 2xy$
16.  $f(x, y) = 3 + 2x + 2y - 2x^2 - 2xy - y^2$
17.  $f(x, y) = x^3 - y^3 - 2xy + 6$
18.  $f(x, y) = x^3 + 3xy + y^3$
19.  $f(x, y) = 6x^2 - 2x^3 + 3y^2 + 6xy$
20.  $f(x, y) = 3y^2 - 2y^3 - 3x^2 + 6xy$

21.  $f(x, y) = 9x^3 + y^3/3 - 4xy$   
 22.  $f(x, y) = 8x^3 + y^3 + 6xy$   
 23.  $f(x, y) = x^3 + y^3 + 3x^2 - 3y^2 - 8$   
 24.  $f(x, y) = 2x^3 + 2y^3 - 9x^2 + 3y^2 - 12y$   
 25.  $f(x, y) = 4xy - x^4 - y^4$   
 26.  $f(x, y) = x^4 + y^4 + 4xy$   
 27.  $f(x, y) = \frac{1}{x^2 + y^2 - 1}$   
 28.  $f(x, y) = \frac{1}{x} + xy + \frac{1}{y}$   
 29.  $f(x, y) = y \sin x$   
 30.  $f(x, y) = e^{2x} \cos y$

### Finding Absolute Extrema

In Exercises 31–38, find the absolute maxima and minima of the functions on the given domains.

31.  $f(x, y) = 2x^2 - 4x + y^2 - 4y + 1$  on the closed triangular plate bounded by the lines  $x = 0$ ,  $y = 2$ ,  $y = 2x$  in the first quadrant  
 32.  $D(x, y) = x^2 - xy + y^2 + 1$  on the closed triangular plate in the first quadrant bounded by the lines  $x = 0$ ,  $y = 4$ ,  $y = x$   
 33.  $f(x, y) = x^2 + y^2$  on the closed triangular plate bounded by the lines  $x = 0$ ,  $y = 0$ ,  $y + 2x = 2$  in the first quadrant  
 34.  $T(x, y) = x^2 + xy + y^2 - 6x$  on the rectangular plate  $0 \leq x \leq 5$ ,  $-3 \leq y \leq 3$   
 35.  $T(x, y) = x^2 + xy + y^2 - 6x + 2$  on the rectangular plate  $0 \leq x \leq 5$ ,  $-3 \leq y \leq 0$   
 36.  $f(x, y) = 48xy - 32x^3 - 24y^2$  on the rectangular plate  $0 \leq x \leq 1$ ,  $0 \leq y \leq 1$   
 37.  $f(x, y) = (4x - x^2) \cos y$  on the rectangular plate  $1 \leq x \leq 3$ ,  $-\pi/4 \leq y \leq \pi/4$  (Fig. 12.53)



12.53 The function and domain in Exercise 37.

38.  $f(x, y) = 4x - 8xy + 2y + 1$  on the triangular plate bounded by the lines  $x = 0$ ,  $y = 0$ ,  $x + y = 1$  in the first quadrant  
 39. Find two numbers  $a$  and  $b$  with  $a \leq b$  such that

$$\int_a^b (6 - x - x^2) dx$$

has its largest value.

40. Find two numbers  $a$  and  $b$  with  $a \leq b$  such that


$$\int_a^b (24 - 2x - x^2)^{1/3} dx$$

has its largest value.

41. *Temperatures.* The flat circular plate in Fig. 12.54 has the shape of the region  $x^2 + y^2 \leq 1$ . The plate, including the boundary where  $x^2 + y^2 = 1$ , is heated so that the temperature at the point  $(x, y)$  is

$$T(x, y) = x^2 + 2y^2 - x.$$

Find the temperatures at the hottest and coldest points on the plate.




12.54 Curves of constant temperature are called isotherms. The figure shows isotherms of the temperature function  $T(x, y) = x^2 + 2y^2 - x$  on the disk  $x^2 + y^2 \leq 1$  in the  $xy$ -plane. Exercise 41 asks you to locate the extreme temperatures.

42. Find the critical point of

$$f(x, y) = xy + 2x - \ln x^2 y$$

in the open first quadrant ( $x > 0$ ,  $y > 0$ ) and show that  $f$  takes on a minimum there (Fig. 12.55).



12.55 The function  $f(x, y) = xy + 2x - \ln x^2 y$  (selected level curves shown here) takes on a minimum value somewhere in the open first quadrant  $x > 0$ ,  $y > 0$  (Exercise 42).

## Theory and Examples

43. Find the maxima, minima, and saddle points of  $f(x, y)$ , if any, given that

a)  $f_x = 2x - 4y$  and  $f_y = 2y - 4x$   
 b)  $f_x = 2x - 2$  and  $f_y = 2y - 4$   
 c)  $f_x = 9x^2 - 9$  and  $f_y = 2y + 4$

Describe your reasoning in each case.

44. The discriminant  $f_{xx}f_{yy} - f_{xy}^2$  is zero at the origin for each of the following functions, so the second derivative test fails there. Determine whether the function has a maximum, a minimum, or neither at the origin by imagining what the surface  $z = f(x, y)$  looks like. Describe your reasoning in each case.

a)  $f(x, y) = x^2y^2$       b)  $f(x, y) = 1 - x^2y^2$   
 c)  $f(x, y) = xy^2$       d)  $f(x, y) = x^3y^2$   
 e)  $f(x, y) = x^3y^3$       f)  $f(x, y) = x^4y^4$

45. Show that  $(0, 0)$  is a critical point of  $f(x, y) = x^2 + kxy + y^2$  no matter what value the constant  $k$  has. (Hint: Consider two cases:  $k = 0$  and  $k \neq 0$ .)

46. For what values of the constant  $k$  does the second derivative test guarantee that  $f(x, y) = x^2 + kxy + y^2$  will have a saddle point at  $(0, 0)$ ? a local minimum at  $(0, 0)$ ? For what values of  $k$  is the second derivative test inconclusive? Give reasons for your answers.

47. a) If  $f_x(a, b) = f_y(a, b) = 0$ , must  $f$  have a local maximum or minimum value at  $(a, b)$ ? Give reasons for your answer.  
 b) Can you conclude anything about  $f(a, b)$  if  $f$  and its first and second partial derivatives are continuous throughout a disk centered at  $(a, b)$  and  $f_{xx}(a, b)$  and  $f_{yy}(a, b)$  differ in sign? Give reasons for your answer.

48. Using the proof of Theorem 7 given in the text for the case in which  $f$  has a local maximum at  $(a, b)$ , prove the theorem for the case in which  $f$  has a local minimum at  $(a, b)$ .

49. Among all the points on the graph of  $z = 10 - x^2 - y^2$  that lie above the plane  $x + 2y + 3z = 0$ , find the point farthest from the plane.

50. Find the point on the graph of  $z = x^2 + y^2 + 10$  nearest the plane  $x + 2y - z = 0$ .

51. The function  $f(x, y) = x + y$  fails to have an absolute maximum value in the closed first quadrant  $x \geq 0$  and  $y \geq 0$ . Does this contradict the discussion on finding absolute extrema given in the text? Give reasons for your answer.

52. Consider the function  $f(x, y) = x^2 + y^2 + 2xy - x - y + 1$  over the square  $0 \leq x \leq 1$  and  $0 \leq y \leq 1$ .

- a) Show that  $f$  has an absolute minimum along the line segment  $2x + 2y = 1$  in this square. What is the absolute minimum value?  
 b) Find the absolute maximum value of  $f$  over the square.

## Extreme Values on Parametrized Curves

To find the extreme values of a function  $f(x, y)$  on a curve  $x =$

$x(t), y = y(t)$ , we treat  $f$  as a function of the single variable  $t$  and use the Chain Rule to find where  $df/dt$  is zero. As in any other single-variable case, the extreme values of  $f$  are then found among the values at the

- a) critical points (points where  $df/dt$  is zero or fails to exist), and  
 b) endpoints of the parameter domain.

Find the absolute maximum and minimum values of the following functions on the given curves.

53. Functions:

a)  $f(x, y) = x + y$       b)  $g(x, y) = xy$   
 c)  $h(x, y) = 2x^2 + y^2$

Curves:

- i) The semicircle  $x^2 + y^2 = 4, y \geq 0$   
 ii) The quarter circle  $x^2 + y^2 = 4, x \geq 0, y \geq 0$

Use the parametric equations  $x = 2 \cos t, y = 2 \sin t$ .

54. Functions:

a)  $f(x, y) = 2x + 3y$       b)  $g(x, y) = xy$   
 c)  $h(x, y) = x^2 + 3y^2$

Curves:

- i) The semi-ellipse  $(x^2/9) + (y^2/4) = 1, y \geq 0$   
 ii) The quarter ellipse  $(x^2/9) + (y^2/4) = 1, x \geq 0, y \geq 0$

Use the parametric equations  $x = 3 \cos t, y = 2 \sin t$ .

55. Function:  $f(x, y) = xy$

Curves:

- i) The line  $x = 2t, y = t + 1$   
 ii) The line segment  $x = 2t, y = t + 1, -1 \leq t \leq 0$   
 iii) The line segment  $x = 2t, y = t + 1, 0 \leq t \leq 1$

56. Functions:

a)  $f(x, y) = x^2 + y^2$       b)  $g(x, y) = 1/(x^2 + y^2)$

Curves:

- i) The line  $x = t, y = 2 - 2t$   
 ii) The line segment  $x = t, y = 2 - 2t, 0 \leq t \leq 1$

## Least Squares and Regression Lines


When we try to fit a line  $y = mx + b$  to a set of numerical data points  $(x_1, y_1), (x_2, y_2), \dots, (x_n, y_n)$  (Fig. 12.56, on the following page), we usually choose the line that minimizes the sum of the squares of the vertical distances from the points to the line. In theory, this means finding the values of  $m$  and  $b$  that minimize the value of the function

$$w = (mx_1 + b - y_1)^2 + \dots + (mx_n + b - y_n)^2. \quad (1)$$

The values of  $m$  and  $b$  that do this are found with the first and second derivative tests to be

$$m = \frac{(\sum x_k)(\sum y_k) - n \sum x_k y_k}{(\sum x_k)^2 - n \sum x_k^2}, \quad (2)$$

$$b = \frac{1}{n} \left( \sum y_k - m \sum x_k \right), \quad (3)$$



12.56 To fit a line to noncollinear points, we choose the line that minimizes the sum of the squares of the deviations.

with all sums running from  $k = 1$  to  $k = n$ . Many scientific calculators have these formulas built in, enabling you to find  $m$  and  $b$  with only a few key presses after you have entered the data.

The line  $y = mx + b$  determined by these values of  $m$  and  $b$  is called the **least squares line**, **regression line**, or **trend line** for the data under study. Finding a least squares line lets you

1. summarize data with a simple expression,
2. predict values of  $y$  for other, experimentally untried values of  $x$ ,
3. handle data analytically.

**EXAMPLE** Find the least squares line for the points  $(0, 1)$ ,  $(1, 3)$ ,  $(2, 2)$ ,  $(3, 4)$ ,  $(4, 5)$ .

**Solution** We organize the calculations in a table:

$k$	$x_k$	$y_k$	$x_k^2$	$x_k y_k$
1	0	1	0	0
2	1	3	1	3
3	2	2	4	4
4	3	4	9	12
5	4	5	16	20
$\Sigma$	10	15	30	39

Then we find

$$m = \frac{(10)(15) - 5(39)}{(10)^2 - 5(30)} = 0.9 \quad \text{Eq. (2) with } n = 5 \text{ and data from the table}$$


and use the value of  $m$  to find

$$b = \frac{1}{5}(15 - (0.9)(10)) = 1.2. \quad \text{Eq. (3) with } n = 5, m = 0.9$$

The least squares line is  $y = 0.9x + 1.2$  (Fig. 12.57).  $\square$

In Exercises 57–60, use Eqs. (2) and (3) to find the least squares line for each set of data points. Then use the linear equation you obtain to predict the value of  $y$  that would correspond to  $x = 4$ .

57.  $(-1, 2)$ ,  $(0, 1)$ ,  $(3, -4)$       58.  $(-2, 0)$ ,  $(0, 2)$ ,  $(2, 3)$   
59.  $(0, 0)$ ,  $(1, 2)$ ,  $(2, 3)$       60.  $(0, 1)$ ,  $(2, 2)$ ,  $(3, 2)$



12.57 The least squares line for the data in the example.

61. Write a linear equation for the effect of irrigation on the yield of alfalfa by fitting a least squares line to the data in Table 12.1 (from the University of California Experimental Station, *Bulletin* No. 450, p. 8). Plot the data and draw the line.

Table 12.1 Growth of alfalfa

$x$ (total seasonal depth of water applied, in.)	$y$ (average alfalfa yield, tons/acre)
12	5.27
18	5.68
24	6.25
30	7.21
36	8.20
42	8.71

62. *Craters of Mars.* One theory of crater formation suggests that the frequency of large craters should fall off as the square of the diameter (Marcus, *Science*, June 21, 1968, p. 1334). Pictures from *Mariner IV* show the frequencies listed in Table 12.2. Fit a line of the form  $F = m(1/D^2) + b$  to the data. Plot the data and draw the line.

Table 12.2 Crater sizes on Mars

Diameter in km, $D$	$1/D^2$ (for left value of class interval)	Frequency, $F$
32–45	0.001	51
45–64	0.0005	22
64–90	0.00024	14
90–128	0.000123	4

Table 12.3 Compositions by Mozart

Köchel number, K	Year composed, y
1	1761
75	1771
155	1772
219	1775
271	1777
351	1780
425	1783
503	1786
575	1789
626	1791

63. *Köchel numbers.* In 1862, the German musicologist Ludwig von Köchel made a chronological list of the musical works of Wolfgang Amadeus Mozart. This list is the source of the Köchel numbers, or “K numbers,” that now accompany the titles of Mozart’s pieces (Sinfonia Concertante in E-flat major, K.364, for example). Table 12.3 gives the Köchel numbers and composition dates (y) of ten of Mozart’s works.

- Plot y vs. K to show that y is close to being a linear function of K.
- Find a least squares line  $y = mK + b$  for the data and add the line to your plot in (a).
- K.364 was composed in 1779. What date is predicted by the least squares line?

64. *Submarine sinkings.* The data in Table 12.4 show the results of a historical study of German submarines sunk by the U.S. Navy during 16 consecutive months of World War II. The data given for each month are the number of reported sinkings and the number of actual sinkings. The number of submarines sunk was slightly greater than the Navy’s reports implied. Find a least squares line for estimating the number of actual sinkings from the number of reported sinkings.

Table 12.4 Sinkings of German submarines by U.S. during 16 consecutive months of WWII

Month	Guesses by U.S. (reported sinkings) x	Actual number y
1	3	3
2	2	2
3	4	6
4	2	3
5	5	4
6	5	3
7	9	11
8	12	9
9	8	10
10	13	16
11	14	13
12	3	5
13	4	6
14	13	19
15	10	15
16	16	15
123		140

- Calculate the function’s second partial derivatives and find the discriminant  $f_{xx}f_{yy} - f_{xy}^2$ .
- Using the max-min tests, classify the critical points found in (c). Are your findings consistent with your discussion in (c)?

- $f(x, y) = x^2 + y^3 - 3xy$ ,  $-5 \leq x \leq 5$ ,  $-5 \leq y \leq 5$
- $f(x, y) = x^3 - 3xy^2 + y^2$ ,  $-2 \leq x \leq 2$ ,  $-2 \leq y \leq 2$
- $f(x, y) = x^4 + y^2 - 8x^2 - 6y + 16$ ,  $-3 \leq x \leq 3$ ,  $-6 \leq y \leq 6$
- $f(x, y) = 2x^4 + y^4 - 2x^2 - 2y^2 + 3$ ,  $-3/2 \leq x \leq 3/2$ ,  $-3/2 \leq y \leq 3/2$
- $f(x, y) = 5x^6 + 18x^5 - 30x^4 + 30xy^2 - 120x^3$ ,  $-4 \leq x \leq 3$ ,  $-2 \leq y \leq 2$
- $f(x, y) = \begin{cases} x^5 \ln(x^2 + y^2), & (x, y) \neq (0, 0) \\ 0, & (x, y) = (0, 0) \end{cases}$ ,  $-2 \leq x \leq 2$ ,  $-2 \leq y \leq 2$

### CAS Explorations and Projects

In Exercises 65–70, you will explore functions to identify their local extrema. Use a CAS to perform the following steps:

- Plot the function over the given rectangle.
- Plot some level curves in the rectangle.
- Calculate the function’s first partial derivatives and use the CAS equation solver to find the critical points. How do the critical points relate to the level curves plotted in (b)? Which critical points, if any, appear to give a saddle point? Give reasons for your answer.