

To calculate the remaining partial derivatives, we apply what we know about the dependence and independence of the variables involved. As shown in the diagram (5), the variables x , y , and z are independent and $t = x + y$. Hence,

$$\frac{\partial x}{\partial x} = 1, \quad \frac{\partial y}{\partial x} = 0, \quad \frac{\partial z}{\partial x} = 0, \quad \frac{\partial t}{\partial x} = \frac{\partial}{\partial x}(x + y) = (1 + 0) = 1.$$

We substitute these values into Eq. (7) to find $\partial w/\partial x$:

$$\begin{aligned} \left(\frac{\partial w}{\partial x}\right)_{y,z} &= 2x(1) + 0 - 0 + (\cos t)(1) \\ &= 2x + \cos t \\ &= 2x + \cos(x + y). \end{aligned} \quad \begin{array}{l} \text{In terms of the independent} \\ \text{variables} \end{array}$$

Exercises 12.6

Finding Partial Derivatives with Constrained Variables

In Exercises 1–3, begin by drawing a diagram that shows the relations among the variables.

1. If $w = x^2 + y^2 + z^2$ and $z = x^2 + y^2$, find

a) $\left(\frac{\partial w}{\partial y}\right)_z$ b) $\left(\frac{\partial w}{\partial z}\right)_x$ c) $\left(\frac{\partial w}{\partial z}\right)_y$

2. If $w = x^2 + y - z + \sin t$ and $x + y = t$, find

a) $\left(\frac{\partial w}{\partial y}\right)_{x,z}$ b) $\left(\frac{\partial w}{\partial y}\right)_{z,t}$ c) $\left(\frac{\partial w}{\partial z}\right)_{x,y}$

d) $\left(\frac{\partial w}{\partial z}\right)_{y,t}$ e) $\left(\frac{\partial w}{\partial t}\right)_{x,z}$ f) $\left(\frac{\partial w}{\partial t}\right)_{y,z}$

3. Let $U = f(P, V, T)$ be the internal energy of a gas that obeys the ideal gas law $PV = nRT$ (n and R constant). Find

a) $\left(\frac{\partial U}{\partial P}\right)_V$ b) $\left(\frac{\partial U}{\partial T}\right)_V$

4. Find

a) $\left(\frac{\partial w}{\partial x}\right)_y$ b) $\left(\frac{\partial w}{\partial z}\right)_y$

at the point $(x, y, z) = (0, 1, \pi)$ if

$$w = x^2 + y^2 + z^2 \quad \text{and} \quad y \sin z + z \sin x = 0.$$

5. Find

a) $\left(\frac{\partial w}{\partial y}\right)_x$ b) $\left(\frac{\partial w}{\partial y}\right)_z$

at the point $(w, x, y, z) = (4, 2, 1, -1)$ if

$$w = x^2y^2 + yz - z^3 \quad \text{and} \quad x^2 + y^2 + z^2 = 6.$$

6. Find $\left(\frac{\partial u}{\partial y}\right)_x$ at the point $(u, v) = (\sqrt{2}, 1)$ if $x = u^2 + v^2$ and $y = uv$.

7. Suppose that $x^2 + y^2 = r^2$ and $x = r \cos \theta$, as in polar coordinates. Find

$$\left(\frac{\partial x}{\partial r}\right)_\theta \quad \text{and} \quad \left(\frac{\partial r}{\partial x}\right)_y.$$

8. Suppose that

$$w = x^2 - y^2 + 4z + t \quad \text{and} \quad x + 2z + t = 25.$$

Show that the equations

$$\frac{\partial w}{\partial x} = 2x - 1 \quad \text{and} \quad \frac{\partial w}{\partial x} = 2x - 2$$

each give $\partial w/\partial x$, depending on which variables are chosen to be dependent and which variables are chosen to be independent. Identify the independent variables in each case.

Partial Derivatives without Specific Formulas

9. Establish the fact, widely used in hydrodynamics, that if $f(x, y, z) = 0$, then

$$\left(\frac{\partial x}{\partial y}\right)_z \left(\frac{\partial y}{\partial z}\right)_x \left(\frac{\partial z}{\partial x}\right)_y = -1.$$

(Hint: Express all the derivatives in terms of the formal partial derivatives $\partial f/\partial x$, $\partial f/\partial y$, and $\partial f/\partial z$.)

10. If $z = x + f(u)$, where $u = xy$, show that

$$x \frac{\partial z}{\partial x} - y \frac{\partial z}{\partial y} = x.$$

11. Suppose that the equation $g(x, y, z) = 0$ determines z as a differentiable function of the independent variables x and y and that $g_z \neq 0$. Show that

$$\left(\frac{\partial z}{\partial y}\right)_x = -\frac{\partial g/\partial y}{\partial g/\partial z}.$$

12. Suppose that $f(x, y, z, w) = 0$ and $g(x, y, z, w) = 0$ determine z and w as differentiable functions of the independent variables

x and y , and suppose that

$$\frac{\partial f}{\partial z} \frac{\partial g}{\partial w} - \frac{\partial f}{\partial w} \frac{\partial g}{\partial z} \neq 0.$$

Show that

$$\left(\frac{\partial z}{\partial x} \right)_y = - \frac{\frac{\partial f}{\partial x} \frac{\partial g}{\partial w} - \frac{\partial f}{\partial w} \frac{\partial g}{\partial x}}{\frac{\partial f}{\partial z} \frac{\partial g}{\partial w} - \frac{\partial f}{\partial w} \frac{\partial g}{\partial z}}$$

and

$$\left(\frac{\partial w}{\partial y} \right)_x = - \frac{\frac{\partial f}{\partial z} \frac{\partial g}{\partial y} - \frac{\partial f}{\partial y} \frac{\partial g}{\partial z}}{\frac{\partial f}{\partial z} \frac{\partial g}{\partial w} - \frac{\partial f}{\partial w} \frac{\partial g}{\partial z}}.$$

12.7

Directional Derivatives, Gradient Vectors, and Tangent Planes

We know from Section 12.5 that if $f(x, y)$ is differentiable, then the rate at which f changes with respect to t along a differentiable curve $x = g(t)$, $y = h(t)$ is

$$\frac{df}{dt} = \frac{\partial f}{\partial x} \frac{dx}{dt} + \frac{\partial f}{\partial y} \frac{dy}{dt}.$$

At any point $P_0(x_0, y_0) = P_0(g(t_0), h(t_0))$, this equation gives the rate of change of f with respect to increasing t and therefore depends, among other things, on the direction of motion along the curve. This observation is particularly important when the curve is a straight line and t is the arc length parameter along the line measured from P_0 in the direction of a given unit vector \mathbf{u} . For then df/dt is the rate of change of f with respect to distance in its domain in the direction of \mathbf{u} . By varying \mathbf{u} , we find the rates at which f changes with respect to distance as we move through P_0 in different directions. These “directional derivatives” have useful interpretations in science and engineering as well as in mathematics. This section develops a formula for calculating them and proceeds from there to find equations for tangent planes and normal lines on surfaces in space.

Directional Derivatives in the Plane

Suppose that the function $f(x, y)$ is defined throughout a region R in the xy -plane, that $P_0(x_0, y_0)$ is a point in R , and that $\mathbf{u} = u_1 \mathbf{i} + u_2 \mathbf{j}$ is a unit vector. Then the equations

$$x = x_0 + su_1, \quad y = y_0 + su_2$$


parametrize the line through P_0 parallel to \mathbf{u} . The parameter s measures arc length from P_0 in the direction of \mathbf{u} . We find the rate of change of f at P_0 in the direction of \mathbf{u} by calculating df/ds at P_0 (Fig. 12.33):

Definition

The **derivative of f at $P_0(x_0, y_0)$ in the direction of the unit vector $\mathbf{u} = u_1 \mathbf{i} + u_2 \mathbf{j}$** is the number

$$\left(\frac{df}{ds} \right)_{\mathbf{u}, P_0} = \lim_{s \rightarrow 0} \frac{f(x_0 + su_1, y_0 + su_2) - f(x_0, y_0)}{s}, \quad (1)$$

provided the limit exists.



12.33 The rate of change of f in the direction of \mathbf{u} at a point P_0 is the rate at which f changes along this line at P_0 .